

Address:

Phone#:

Record of Immunization(s) Provided at Clinic

NAME:

DOB:

PARIS Id:

Age:

PHN:

The Immunization(s) checked below were given on

Time:

Immunization	✓	Initial	Next Due	Immunization	✓	Initial	Next Due
Diphth, Pert, Tet, Polio, Hib				Meningococcal Conj C			
Diphth, Pert, Tet, Polio				Meningococcal ACYW 35			
Diphth,Pert,Tet,Polio,Hib,HepB				Hepatitis B			
Tetanus, diphtheria, Pertussis				Hepatitis A			
Tetanus, diphtheria, Polio				Hepatitis A/B			
Tetanus, diphtheria				Pneumococcal 07			
Polio				Pneumococcal 23			
Measles, Mumps, Rubella				HIB			
Varicella				Influenza			
Human Papillomavirus				Other			

Phone

to make an appointment.

Immunization History Chart

Description	Dose	Date (DD/MMM/YYYY)	Age Given	See Notes	Description	Dose	Date (DD/MMM/YYYY)	Age Given	See Notes

Note: ED = Date of dose is unknown and has been estimated.

* = Some or all of this immunization does not meet BC Guidelines. This immunization may need to be repeated.

CARE AFTER IMMUNIZATION

Minor Reactions are expected after most vaccines and may consist of soreness, redness, fever and swelling at the injection site.

Reactions usually last 1-2 days and can be controlled by placing cool cloths on the injection site and giving extra fluids.

After immunization, a child with a temperature over 38°C should receive acetaminophen (Tempra, Tylenol, Panadol, Atasol), to make him/her more comfortable.

Reactions to MMR vaccine could include a slight rash and swollen glands which may appear between 3-20 days following the injection. This usually lasts for 1-2 days.

Reactions to the Varicella vaccine could include developing a small varicella-like rash, 5-26 days following the injection.

If you have any concerns or experience a serious reaction, please consult your family doctor or phone the number located on the top of the page.